Rachel M. Frank, MD

Team Physician, *University of Colorado Athletics*Assistant Professor, Department of Orthopaedic Surgery
University of Colorado School of Medicine
www.RachelFrankMD.com



Physical Therapy Prescription – Cartilage Restoration Patella / Trochlea

Name:	
Procedure: R / L	Date of Surgery:
Frequency: 2-3 times per week for weeks	
PHASE I (Weeks 0 – 6): Period of protection, decrease edema	, activate quadriceps
 extension lag Range of Motion: Continuous Passive Motion (CPM) r CPM Protocol: 1 cycle per minute starting 0-30 Therapeutic Exercises: Weeks 0-2: quad sets, calf pumps, passive leg 	roved; discontinue when able to perform SLR without machine for 6-8 hours/day 0° (weeks 0-2), 0-60° (weeks 2-4), 0-90° (weeks 4-6) I hangs to 45° I hangs to
Phase III (Weeks 8 – 12)	
 Weightbearing: Full Range of Motion: Full, painless Therapeutic Exercises: Advance Phase II, begin close raises), begin stationary bike, begin unilateral stance as Modalities: Per therapist, including electrical stimulation 	ctivities and balance training
Phase IV (Weeks 12 – 24)	

Advance Phase III exercises; focus on core/glutes; advance to elliptical, bike, and pool as tolerated

• Encourage maintenance program

Phase V (>6 months): Gradual return to athletic activity

• Return to sport-specific activity and impact when cleared by MD at 8-9 months postop

Signature:	Date:	