#### Rachel M. Frank. MD

Team Physician, *University of Colorado Athletics*Assistant Professor, Department of Orthopaedic Surgery
University of Colorado School of Medicine
www.RachelFrankMD.com



# **Physical Therapy Prescription – Meniscus Allograft Transplantation**

Name:	Date:
Procedure: R / L meniscus transplantation	Date of Surgery:
Frequency: 2-3 times per week for weeks	

PHASE I (Weeks 0 – 6): Period of protection, decrease edema, activate quadriceps

- Weightbearing: Heel-touch WB with crutches in brace
- Hinged Knee Brace:
  - Weeks 0-2: Locked in full extension for ambulation and sleeping
  - Weeks 2-6: Unlocked 0-90° for ambulation and removed while sleeping
- Range of Motion: AAROM → AROM as tolerated; no weight-bearing with knee flexion angles >90°
- Therapeutic Exercises: Patellar mobs, quad/hamstring sets, heel slides, Gastroc/Soleus stretching, SLRs with brace in full extension until quad strength prevents extension lag, total gym (closed chain)
  - Avoid tibial rotation for first 8 weeks to protect the allograft
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

# Phase II (Weeks 6 - 12)

- Weightbearing: Increase 25% per week, progress to full WBAT by week 8
- Hinged Knee Brace: Wean out of brace between 6-8 weeks: discontinue at 8 weeks
- Range of Motion: Full
- Therapeutic Exercises: Progress Phase I exercises, progress closed chain exercises, begin hamstring exercises, toe lunges 0-90°, leg press 0-90°, begin proprioception exercises, work on balance/core/glutes
  - Begin use of the stationary bike at week 8
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

## Phase III (Weeks 12 - 16)

- Range of Motion: Full, painless
- Therapeutic Exercises: Progress Phase II exercises, advance closed chain strengthening exercises and proprioception activities; focus on single-leg balance; begin elliptical; progress with stationary bike
  - Jogging okay at 12 weeks
  - Swimming okay at 16 weeks
- Modalities: Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

## Phase IV (Weeks 16 - 24)

- Progress Phase III exercises, focus on single-leg strength, progress plyometrics and sport-specific training
- Consider functional sports assessment

#### Phase V (>6 months): Gradual return to athletic activity

- Gradual return to sports participation after completion of functional sports assessment
- Encourage maintenance program based off functional sports assessment

Signature: Date: